

Perinatal Pandemic (H1N1) 2009 Infection, Thailand

To the Editor: Infection with influenza A pandemic (H1N1) 2009 has been reported worldwide following initial identification of the virus in April 2009 (1). The groups at highest risk for infection or influenza-related complications include pregnant women and children (2). We report a case of pandemic (H1N1) 2009 infection in a newborn whose mother became ill with pandemic (H1N1) 2009 during the perinatal period.

A newborn girl showed signs of respiratory distress. The relevant perinatal history was maternal illness with pandemic (H1N1) 2009 7 days before delivery. The infant, who had a birth weight of 1,560 grams, was delivered by emergency cesarean section after the mother experienced cardiopulmonary failure at the gestational age of 31 weeks. Apgar scores were 9 and 9 at 1 and 5 minutes, respectively. Physical examination at birth showed a premature infant girl with mild subcostal retraction. Oxygen saturation at room air was 91%–99%. Other results of the physical examination were unremarkable.

Initial management included routine care for premature infants. On the basis of the perinatal history, a throat swab specimen was collected for pandemic (H1N1) 2009 testing by PCR and oseltamivir, 6 mg, was administered every 12 hours (4 mg/kg/day). The specimen obtained from the throat swab was positive for pandemic (H1N1) 2009 by real-time PCR. The infant required oxygen supplementation. At day 2 of life, acute renal failure with an elevated plasma creatinine level of 1.1 mg/dL developed in the infant. Chest radiograph showed minimal pulmonary infiltrations. She was started on cefotaxime for suspected sepsis. Oseltamivir dosage was adjusted based on the glomerular fil-

tration rate estimated by the formula of Schwartz et al. (3) of 10.5 mL/min/1.73m² to 3 mg every 12 hours to complete 10 doses (2 mg/kg/day).

Infection of the patient was confirmed by real-time reverse transcription-PCR of the throat swab specimen and by a 4× increase in antibodies against the virus by hemagglutination inhibition test (HI). Antibody titers against pandemic influenza (H1N1) 2009 by HI with turkey erythrocytes (4) on days 10, 24, and 42 of life were 10, 160, and 320, respectively (Figure). At day 4 of life, repeated PCR performed on a throat swab specimen was negative for pandemic (H1N1) 2009. Oxygen supplementation was gradually decreased and finally discontinued. Her room air oxygen saturation was 95%–98%. Her clinical symptoms gradually improved. Hemoculture was negative after 72 hours. The antimicrobial drugs were given over an 8-day course. Plasma creatinine decreased to 0.9 mg/dL and 0.6 mg/dL at days 6 and 7 of life, respectively. Her average urine output was 2–3 mL/kg/h. She was discharged at the age of 28 days with a body weight of 2,070 grams.

Pregnant women are one of the highest risk groups for influenza A infection and influenza-associated complications, including increased maternal and perinatal illness and death rates (5). Thus, pregnant women are given first priority to receive influenza vaccination. When influenza develops in mothers during the perinatal period, newborns can be infected transplacentally during maternal viremia or by respiratory droplets after birth. Transplacental infection of influenza A is rare (5), however, and there have been only a few case reports (6,7). Viremia is more frequent and more extensive in pregnant women due to depressed cell-mediated immune response during the pregnancy (8). Our patient was likely infected in utero because she was delivered by cesarean section and was never exposed to her mother, who required intensive cardiopulmonary support at the time of delivery. (The mother died from respiratory failure 7 days after the cesarean section.)

Clinical manifestations in this patient, including respiratory distress and acute renal failure, were nonspe-




Figure. Antibody titer against influenza A pandemic (H1N1) 2009 by hemagglutination inhibition (HI) test on days 10, 24, and 42 of life of the patient. A color version of this figure is available online (www.cdc.gov/EID/content/16/2/343-F.htm).

cific. The high plasma creatinine level in the newborn sometimes reflects the mother's plasma creatinine level (9). However, kidney function of the mother of the newborn was within normal limits at the time of Cesarean section; plasma creatinine level of 0.7 mg/dL. An elevated plasma creatinine level is observed frequently in premature infants due to immaturity of the kidney tissue and will usually decrease within a few weeks. Oseltamivir was administered with dose adjustment based on the infant's estimated glomerular filtration rate. The recommended dose of oseltamivir for glomerular filtration rate <30 mL/min/1.73 m² is 2–3 mg/kg/day, based on preliminary data obtained by a National Institutes of Health–funded Collaborative Antiviral Study Group (10). The success of our management strategy for this case suggests early treatment with oseltamivir can prevent severe illness in newborns with perinatal influenza A pandemic (H1N1) 2009 infection.

Acknowledgments

We thank the staff of Ratchaburi Hospital for taking care of the neonate and Petra Hirsch for reviewing the manuscript.

This research was supported by the Center of Excellence in Clinical Virology Fund, Faculty of Medicine, Chulalongkorn University and Hospital; Thai Red Cross Society; Commission on Higher Education, Ministry of Education, Chulalongkorn University; and Chulalongkorn University Centenary Academic Development Project.

**Wut Dulyachai, Jarika Makkoch,
Pornpimol Rianthavorn,
Mutita Changpinyo,
Slinporn Prayangprecha,
Sunchai Payungporn,
Rachod Tantilertcharoen,
Pravina Kitikoon,
and Yong Poovorawan**

Author affiliations: Ratchaburi Hospital, Ratchaburi, Thailand (W. Dulyachai, M. Changpinyo); Chulalongkorn University, Bangkok, Thailand (J. Makkoch, P. Rianthavorn, S. Prayangprecha, S. Payungporn, R. Tantilertcharoen, P. Kitikoon, Y. Poovorawan); and King Chulalongkorn Memorial Hospital, Bangkok (P. Rianthavorn)

DOI: 10.3201/eid1602.091733

References

- Centers for Disease Control and Prevention. Swine influenza A (H1N1) infection in two children—southern California, March–April 2009. *MMWR Morb Mortal Wkly Rep.* 2009;58:400–42.
- Centers for Disease Control and Prevention. Use of influenza A (H1N1) 2009 monovalent vaccine: recommendations of the Advisory Committee on Immunization Practices (ACIP), 2009. *MMWR Recomm Rep.* 2009;58(RR-10):1–8.
- Schwartz GJ, Brion LP, Spitzer A. The use of plasma creatinine concentration for estimating glomerular filtration rates in infants, children, and adolescents. *Pediatr Clin North Am.* 1987;34:571–90.
- Rowe T, Abernathy RA, Hu-Primmer J, Thompson WW, Lu X, Lim W, et al. Detection of antibody to avian influenza A (H5N1) virus in human serum by using a combination of serologic assays. *J Clin Microbiol.* 1999;37:937–43.
- Irving WL, James DK, Stephenson T, Laing P, Jameson C, Oxford JS, et al. Influenza virus infection in the second and third trimesters of pregnancy: a clinical and seroepidemiological study. *BJOG.* 2000;107:1282–9. DOI: 10.1111/j.1471-0528.2000.tb11621.x
- McGregor JA, Burns JC, Levin MJ, Burlington B, Meiklejohn G. Transplacental passage of influenza A/Bangkok (H3N2) mimicking amniotic fluid infection syndrome. *Am J Obstet Gynecol.* 1984;149:856–9.
- Yawn DH, Pyeatte JC, Joseph JM, Eichler SL, Garcia-Bunuel R. Transplacental transfer of influenza virus. *JAMA.* 1971;216:1022–3. DOI: 10.1001/jama.216.6.1022
- Purtilo DT, Hallgren HM, Yunis EJ. Depressed maternal lymphocyte response to phytohemagglutinin in human pregnancy. *Lancet.* 1972;1:769. DOI: 10.1016/S0140-6736(72)90522-3
- Bueva A, Guignard JP. Renal function in preterm infants. *Pediatr Res.* 1994;36:572–7.
- Allen U, Blumberg EA, Fischer SA, Green M, Humar A, Ison MG, et al. American Society of Transplantation Infectious Diseases Community of Practice Transplant Infectious Disease Section of the Transplantation Society Guidance on Novel Influenza A/H1N1 [cited 2009 Nov 19]. <http://www.transplantation-soc.org/downloads>

Address for correspondence: Yong Poovorawan, Center of Excellence in Clinical Virology, Department of Pediatrics, Faculty of Medicine, Chulalongkorn University, Bangkok, 10330 Thailand; email: yong.p@chula.ac.th

Bronchial Casts and Pandemic (H1N1) 2009 Virus Infection

To the Editor: In the late 1990s, triple-reassortant influenza A viruses containing genes from avian, human, and swine influenza viruses emerged and became enzootic in swine herds in North America (1). The first 11 human cases of novel influenza A virus infection were reported to the Centers for Disease Control and Prevention (CDC; Atlanta, GA, USA) from December 2005 through February 2009 (1). In response to those reports, surveillance for human infection with nonsubtypeable influenza A viruses was implemented.

In the spring of 2009, outbreaks of febrile respiratory infections caused by a novel influenza A virus (H1N1) were reported among persons in Mexico, the United States, and Canada (2). Patient specimens were sent to CDC for real-time reverse transcription–PCR (RT-PCR) testing, and from April 15 through May 5, 2009, a total of 642 infections with the virus, now called pandemic (H1N1) 2009 virus, were confirmed. Of those 642 patients, 60% were ≤18 years of age, indicating that